

PCT/EP 03 / 001 53

REC'D 14 FEB 2003

WIPO

PCT

PA 916063

THE UNITED STATES OF AMERICA

TO ALL TO WHOM THESE PRESENTS SHALL COME:

UNITED STATES DEPARTMENT OF COMMERCE

United States Patent and Trademark Office

October 29, 2002

THIS IS TO CERTIFY THAT ANNEXED HERETO IS A TRUE COPY FROM THE RECORDS OF THE UNITED STATES PATENT AND TRADEMARK OFFICE OF THOSE PAPERS OF THE BELOW IDENTIFIED PATENT APPLICATION THAT MET THE REQUIREMENTS TO BE GRANTED A FILING DATE UNDER 35 USC 111.

APPLICATION NUMBER: 60/347,264

FILING DATE: January 10, 2002

**PRIORITY
DOCUMENT**

SUBMITTED OR TRANSMITTED IN
COMPLIANCE WITH RULE 17.1(a) OR (b)

**By Authority of the
COMMISSIONER OF PATENTS AND TRADEMARKS**



T. Wallace
T. WALLACE
Certifying Officer

01/10/02

JC928 U.S. PTO



Docket Number 4-32304P1/PROV

FILING BY "EXPRESS MAIL" UNDER 37 CFR 1.10

EL790911262US
Express Mail Label NumberJanuary 10, 2002
Date of DepositJC973 U.S. PTO
60/347264Address to: Assistant Commissioner for Patents
Box Provisional Patent Application
Washington, DC 20231

PROVISIONAL APPLICATION FOR PATENT COVER SHEET

Transmitted herewith for filing under 37 CFR §1.53(c) is the PROVISIONAL APPLICATION for patent of

INVENTOR(S)		
Given Name (first and middle [if any])	Family Name or Surname	Residence (City and either State or Foreign Country)
Margaret Forney	Prescott	Millburn, New Jersey
TITLE OF THE INVENTION (280 characters max) ORGANIC COMPOUNDS		
CORRESPONDENCE ADDRESS Direct all correspondence to the address associated with Customer No. 001095, which is currently: Thomas Hoxie Novartis Corporation Patent and Trademark Dept. 564 Morris Avenue Summit, NJ 07901-1027		
ENCLOSED APPLICATION PARTS (check all that apply) <input checked="" type="checkbox"/> Specification (Including Any Claims and Abstract) - 17 pages <input type="checkbox"/> Drawings - sheets <input checked="" type="checkbox"/> Other (specify): Application Data Sheet		
METHOD OF PAYMENT The Commissioner is hereby authorized to charge filing fee and any additional fees required to Deposit Account Number: 19-0134 in the name of Novartis Corporation.		
PROVISIONAL FILING FEE AMOUNT: \$ 160		

☐ U.S. Government agency and contract number: (if the invention was made by an agency of the United States Government or under a contract with an agency of the United States Government.)

Respectfully submitted,

Date: January 10, 2002

Thomas R. Savitsky
Attorney for Applicant
Reg. No. 31,661
Tel. No. (908) 522-6765

INVENTOR INFORMATION

Inventor One Given Name:: Margaret Rorney
Family Name:: Prescott
Postal Address Line One:: 2 Farview Road
City:: Millburn
State or Province:: New Jersey
Country:: US
Postal or Zip Code:: 07041
City of Residence:: Millburn
State or Province of Residence:: New Jersey
Country of Residence:: US
Citizenship Country:: United States

CORRESPONDENCE INFORMATION

Correspondence Customer Number:: 001095
Fax One:: 908-522-6955

APPLICATION INFORMATION

Title Line One:: ORGANIC COMPOUNDS
Formal Drawings?: No
Application Type:: Provisional
Docket Number:: 4-32304P1
Secrecy Order in Parent Appl.?: No

REPRESENTATIVE INFORMATION

Representative Customer Number:: 1095
Source:: PrintEFS Version 1.0.1

2007-01-10 10:24:04

Organic Compounds

The present invention relates to drug delivery systems for the prevention and treatment of proliferative diseases, particularly vascular diseases.

Many humans suffer from circulatory diseases caused by a progressive blockage of the blood vessels that perfuse the heart and other major organs. Severe blockage of blood vessels in such humans often leads to ischemic injury, hypertension, stroke or myocardial infarction. Atherosclerotic lesions which limit or obstruct coronary or periphery blood flow are the major cause of ischemic disease related morbidity and mortality including coronary heart disease and stroke. To stop the disease process and prevent the more advanced disease states in which the cardiac muscle or other organs are compromised, medical revascularization procedures such as percutaneous transluminal coronary angioplasty (PCTA), percutaneous transluminal angioplasty (PTA), atherectomy, bypass grafting or other types of vascular grafting procedures are used.


Re-narrowing (restenosis) of an atherosclerotic coronary artery after various revascularization procedures occurs in 10-80% of patients undergoing this treatment, depending on the procedure used and the arterial site. Besides opening an artery obstructed by atherosclerosis, revascularization also injures endothelial cells and smooth muscle cells within the vessel wall, thus initiating a thrombotic and inflammatory response. Cell derived growth factors such as platelet derived growth factor, infiltrating macrophages, leukocytes or the smooth muscle cells themselves provoke proliferative and migratory responses in the smooth muscle cells. Simultaneous with local proliferation and migration, inflammatory cells also invade the site of vascular injury and may migrate to the deeper layers of the vessel wall. Proliferation/migration usually begins within one to two days post-injury and, depending on the revascularization procedure used, continues for days and weeks.

Both cells within the atherosclerotic lesion and those within the media migrate, proliferate and/or secrete significant amounts of extracellular matrix proteins. Proliferation, migration and extracellular matrix synthesis continue until the damaged endothelial layer is repaired at which time proliferation slows within the intima. The newly formed tissue is called neointima, intimal thickening or restenotic lesion and usually results in narrowing of the vessel lumen. Further lumen narrowing may take place due to constructive remodeling, e.g. vascular remodeling, leading to further intimal thickening or hyperplasia.

2007-07-24 10:01:00

It has now been found that rapamycin derivatives having mTOR inhibiting properties, optionally in conjunction with other active compounds, e.g. antiproliferative compounds, have beneficial effects when locally applied to the lesions sites.

1001 1002 1003 1004 1005 1006 1007 1008 1009 1010 1011 1012 1013 1014 1015 1016 1017 1018 1019 1020 1021 1022 1023 1024 1025 1026 1027 1028 1029 1030 1031 1032 1033 1034 1035 1036 1037 1038 1039 1040 1041 1042 1043 1044 1045 1046 1047 1048 1049 1050 1051 1052 1053 1054 1055 1056 1057 1058 1059 1060 1061 1062 1063 1064 1065 1066 1067 1068 1069 1070 1071 1072 1073 1074 1075 1076 1077 1078 1079 1080 1081 1082 1083 1084 1085 1086 1087 1088 1089 1090 1091 1092 1093 1094 1095 1096 1097 1098 1099 1100 1101 1102 1103 1104 1105 1106 1107 1108 1109 1110 1111 1112 1113 1114 1115 1116 1117 1118 1119 1120 1121 1122 1123 1124 1125 1126 1127 1128 1129 1130 1131 1132 1133 1134 1135 1136 1137 1138 1139 1140 1141 1142 1143 1144 1145 1146 1147 1148 1149 1150 1151 1152 1153 1154 1155 1156 1157 1158 1159 1160 1161 1162 1163 1164 1165 1166 1167 1168 1169 1170 1171 1172 1173 1174 1175 1176 1177 1178 1179 1180 1181 1182 1183 1184 1185 1186 1187 1188 1189 1190 1191 1192 1193 1194 1195 1196 1197 1198 1199 1200 1201 1202 1203 1204 1205 1206 1207 1208 1209 1210 1211 1212 1213 1214 1215 1216 1217 1218 1219 1220 1221 1222 1223 1224 1225 1226 1227 1228 1229 1230 1231 1232 1233 1234 1235 1236 1237 1238 1239 1240 1241 1242 1243 1244 1245 1246 1247 1248 1249 1250 1251 1252 1253 1254 1255 1256 1257 1258 1259 1260 1261 1262 1263 1264 1265 1266 1267 1268 1269 1270 1271 1272 1273 1274 1275 1276 1277 1278 1279 1280 1281 1282 1283 1284 1285 1286 1287 1288 1289 1290 1291 1292 1293 1294 1295 1296 1297 1298 1299 1300 1301 1302 1303 1304 1305 1306 1307 1308 1309 1310 1311 1312 1313 1314 1315 1316 1317 1318 1319 1320 1321 1322 1323 1324 1325 1326 1327 1328 1329 1330 1331 1332 1333 1334 1335 1336 1337 1338 1339 1340 1341 1342 1343 1344 1345 1346 1347 1348 1349 1350 1351 1352 1353 1354 1355 1356 1357 1358 1359 1360 1361 1362 1363 1364 1365 1366 1367 1368 1369 1370 1371 1372 1373 1374 1375 1376 1377 1378 1379 1380 1381 1382 1383 1384 1385 1386 1387 1388 1389 1390 1391 1392 1393 1394 1395 1396 1397 1398 1399 1400 1401 1402 1403 1404 1405 1406 1407 1408 1409 1410 1411 1412 1413 1414 1415 1416 1417 1418 1419 1420 1421 1422 1423 1424 1425 1426 1427 1428 1429 1430 1431 1432 1433 1434 1435 1436 1437 1438 1439 1440 1441 1442 1443 1444 1445 1446 1447 1448 1449 1450 1451 1452 1453 1454 1455 1456 1457 1458 1459 1460 1461 1462 1463 1464 1465 1466 1467 1468 1469 1470 1471 1472 1473 1474 1475 1476 1477 1478 1479 1480 1481 1482 1483 1484 1485 1486 1487 1488 1489 1490 1491 1492 1493 1494 1495 1496 1497 1498 1499 1500 1501 1502 1503 1504 1505 1506 1507 1508 1509 1510 1511 1512 1513 1514 1515 1516 1517 1518 1519 1520 1521 1522 1523 1524 1525 1526 1527 1528 1529 1530 1531 1532 1533 1534 1535 1536 1537 1538 1539 1540 1541 1542 1543 1544 1545 1546 1547 1548 1549 1550 1551 1552 1553 1554 1555 1556 1557 1558 1559 1560 1561 1562 1563 1564 1565 1566 1567 1568 1569 1570 1571 1572 1573 1574 1575 1576 1577 1578 1579 1580 1581 1582 1583 1584 1585 1586 1587 1588 1589 1590 1591 1592 1593 1594 1595 1596 1597 1598 1599 1600 1601 1602 1603 1604 1605 1606 1607 1608 1609 1610 1611 1612 1613 1614 1615 1616 1617 1618 1619 1620 1621 1622 1623 1624 1625 1626 1627 1628 1629 1630 1631 1632 1633 1634 1635 1636 1637 1638 1639 1640 1641 1642 1643 1644 1645 1646 1647 1648 1649 1650 1651 1652 1653 1654 1655 1656 1657 1658 1659 1660 1661 1662 1663 1664 1665 1666 1667 1668 1669 1670 1671 1672 1673 1674 1675 1676 1677 1678 1679 1680 1681 1682 1683 1684 1685 1686 1687 1688 1689 1690 1691 1692 1693 1694 1695 1696 1697 1698 1699 1700 1701 1702 1703 1704 1705 1706 1707 1708 1709 1710 1711 1712 1713 1714 1715 1716 1717 1718 1719 1720 1721 1722 1723 1724 1725 1726 1727 1728 1729 1730 1731 1732 1733 1734 1735 1736 1737 1738 1739 1740 1741 1742 1743 1744 1745 1746 1747 1748 1749 1750 1751 1752 1753 1754 1755 1756 1757 1758 1759 1760 1761 1762 1763 1764 1765 1766 1767 1768 1769 1770 1771 1772 1773 1774 1775 1776 1777 1778 1779 1780 1781 1782 1783 1784 1785 1786 1787 1788 1789 1790 1791 1792 1793 1794 1795 1796 1797 1798 1799 1800 1801 1802 1803 1804 1805 1806 1807 1808 1809 1810 1811 1812 1813 1814 1815 1816 1817 1818 1



R₁ is CH₃ or C₃₋₆alkynyl,

X is =O, (H,H) or (H,OH)

A preferred compound is e.g. 40-0-(2-hydroxyethyl)-rapamycin disclosed in Example 8 in WO 94/09010, or 32-deoxorapamycin as disclosed in Example 1 in WO 96/41807.

According to the invention, the rapamycin derivative may be applied as the sole active ingredient or in conjunction with an immunosuppressive agent, e.g. a calcineurin inhibitor, e.g. a cyclosporin, for example cyclosporin A, or FK506, an EDG-Receptor agonist, e.g. FTY720, an anti-inflammatory agent, e.g. a steroid, e.g. a corticosteroid, e.g. dexamethasone or prednisone, a NSAID, e.g. a cyclooxygenase inhibitor, e.g. a cox-2 inhibitor, e.g. celecoxib, rofecoxib, etoricoxib or valdecoxib, or an ascomycin, e.g. ASM981,

an anti-thrombotic or anti-coagulant agent, e.g. heparin, a IIb/IIIa inhibitor, etc. an antiproliferative agent, e.g. a microtubule stabilizing or destabilizing agent including but not limited to taxanes, e.g. taxol, paclitaxel or docetaxel, vinca alkaloids, e.g. vinblastine, especially vinblastine sulfate, vincristine especially vincristine sulfate, and vinorelbine, discodermolides or epothilones or a derivative thereof, e.g. epothilone B or a derivative thereof, a tyrosine kinase inhibitor, e.g. staurosporin and related small molecules, e.g. UCN-01, BAY 43-9006, Bryostatin 1, Perifosine, Limofosine, midostaurin, RO318220, RO320432, GO 6976, Isis 3521, LY333531, LY379196, SU5416, SU6668, AG1296 etc., a compound or antibody which inhibits the PDGF receptor tyrosine kinase or a compound which binds to PDGF or reduces expression of the PDGF receptor e.g. STI571, CT52923, RP-1776, GFB-111, pyrrolo[3,4-c]-beta-carboline-diones, etc., a compound or antibody which inhibits the EGF receptor tyrosine kinase or a compound which binds to EGF or reduces expression of the EGF receptor e.g. the compounds disclosed in WO97/02266, e.g. the compound of example 39, retinoic acid, ZD1839 (Iressa), alpha-, gamma- or delta-tocopherol or alpha-, gamma- or delta-tocotrienol, or compounds affecting GRB2, IMC-C225, a compound or antibody which inhibits the VEGF receptor tyrosine kinase or a VEGF receptor or a compound which binds to VEGF, e.g. proteins, small molecules or monoclonal antibodies generically and specifically disclosed in WO 98/35958, e.g. 1-(4-chloroanilino)-4-(4-pyridylmethyl)phthalazine or a pharmaceutically acceptable salt thereof, e.g. the succinate, or in WO 00/09495, WO 00/27820, WO 00/59509, WO 98/11223, WO 00/27819, WO 00/37502, WO 94/10202 and EP 0 769 947, those as described by M. Prewett et al in Cancer Research 59 (1999) 5209-5218, by F. Yuan et al in Proc. Natl. Acad. Sci. USA, vol. 93, pp. 14765-14770, Dec. 1996, by Z. Zhu et al in Cancer Res. 58, 1998, 3209-3214, by J. Mordenti et al in Toxicologic Pathology, Vol. 27, no. 1, pp 14-21, 1999, AngiostatinTM, described by M. S. O'Reilly et al, Cell 79, 1994, 315-328, EndostatinTM, described by M. S. O'Reilly et al, Cell 88, 1997, 277-285, anthranilic acid amides, ZD4190; ZD6474, SU5416, SU6668 or anti-VEGF antibodies or anti-VEGF receptor antibodies, e.g. RhuMab, a statin, e.g. having HMG-CoA reductase inhibition activity, e.g. fluvastatin, lovastatin, simvastatin, pravastatin, atorvastatin, cerivastatin, pitavastatin, rosuvastatin or nivastatin, a compound, protein, growth factor or compound stimulating growth factor production that will enhance endothelial regrowth of the luminal endothelium, e.g. FGF, IGF, a matrix metalloproteinase inhibitor, e.g. batimistat, marimistat, trocade, CGS 27023, RS 130830 or AG3340, a modulator (i.e. antagonists or agonists) of kinases, e.g. JNK, ERK1/2, MAPK or STAT, or a

60347264-01102

compound stimulating the release of (NO) or a NO donor, e.g. diazeniumdiolates, S-nitrosothiols, mesoionic oxatriazoles, a combination of isosorbide.

The present invention also provides the local administration or delivery of rapamycin in conjunction with a calcineurin inhibitor, e.g. as disclosed above, an EDG-Receptor agonist, e.g. as disclosed above, a microtubule stabilizing or destabilizing agent, e.g. as disclosed above, a compound or antibody which inhibits the PDGF receptor tyrosine kinase or a compound which binds to PDGF or reduces expression of the PDGF receptor, e.g. as disclosed above, a compound or antibody which inhibits the EGF receptor tyrosine kinase or a compound which binds to EGF or reduces expression of the EGF receptor, e.g. as disclosed above, a compound or antibody which inhibits the VEGF receptor tyrosine kinase or a VEGF receptor or a compound which binds to VEGF, e.g. as disclosed above, a statin, e.g. as disclosed above, a compound, protein, growth factor or compound stimulating growth factor production that will enhance endothelial regrowth of the luminal endothelium, e.g. as disclosed above, a matrix metalloproteinase inhibitor, e.g. as disclosed above, an inhibitor of a modulator (i.e. antagonists or agonists) of kinases, e.g. as disclosed above, or a compound stimulating the release of (NO) or a NO donor, e.g. as disclosed above.

In accordance with the particular findings of the present invention, there is provided

1. A method for preventing or treating smooth muscle cell proliferation and migration in hollow tubes, or increased cell proliferation or decreased apoptosis or increased matrix deposition in a mammal in need thereof, comprising local administration of a therapeutically effective amount of a rapamycin derivative having mTOR inhibiting properties, optionally in conjunction with one or more other active ingredients, e.g. as disclosed above, or a therapeutically effective amount of rapamycin in conjunction with one or more other active ingredients as disclosed above.
2. A method for the treatment of intimal thickening in vessel walls comprising the controlled delivery from any catheter-based device or intraluminal medical device of a therapeutically effective amount of a rapamycin derivative having mTOR inhibiting properties, optionally in conjunction with one or more other active ingredients, e.g. as disclosed above, or a therapeutically effective amount of rapamycin in conjunction with one or more other active ingredients as disclosed above.

Preferably the treatment of intimal thickening in vessel walls is stenosis, restenosis, e.g. following revascularization or neovascularization, and/or inflammation and/or thrombosis.

2025-07-24 10:10:02

3. A drug delivery device or system comprising a) a medical device adapted for local application or administration in hollow tubes, e.g. a catheter-based delivery device or intraluminal medical device, and b) a therapeutic dosage of a rapamycin derivative having mTOR inhibiting properties or rapamycin, optionally in conjunction with a therapeutic dosage of one or more other active ingredients, e.g. as disclosed above, each being releasably affixed to the catheter-based delivery device or medical device.

Such a local delivery device or system can be used to reduce stenosis or restenosis as an adjunct to revascularization, bypass or grafting procedures performed in any vascular location including coronary arteries, carotid arteries, renal arteries, peripheral arteries, cerebral arteries or any other arterial or venous location, to reduce anastomotic stenosis such as in the case of arterial-venous dialysis access with or without polytetrafluoroethylene grafting and with or without stenting, or in conjunction with any other heart or transplantation procedures, or congenital vascular interventions.

In a further embodiment, the present invention also provides a drug delivery system or device as disclosed above comprising b) a therapeutic dosage of a compound or antibody which inhibits the PDGF receptor tyrosine kinase or a compound which binds to PDGF or reduces expression of the PDGF receptor e.g. as disclosed above, a compound or antibody which inhibits the EGF receptor tyrosine kinase or a compound which binds to EGF or reduces expression of the EGF receptor e.g. as disclosed above, a compound or antibody which inhibits the VEGF receptor tyrosine kinase or a VEGF receptor or a compound which binds to VEGF, e.g. as disclosed above, each being releasably affixed to the catheter-based delivery device or medical device.

Rapamycin or rapamycin derivative having mTOR inhibiting properties will be referred to hereinafter as "drug". The other active ingredients which may be used in conjunction with rapamycin or a rapamycin derivative, e.g. as disclosed above, will be referred to hereinafter collectively as "adjunct". Drug(s) shall mean drug or drug+adjunct.

The local administration preferably takes place at or near the vascular lesions sites.

The administration may be by one or more of the following routes: via catheter or other intravascular delivery system, intranasally, intrabronchially, interperitoneally or esophageal. Hollow tubes include circulatory system vessels such as blood vessels (arteries or veins), tissue lumen, lymphatic pathways, digestive tract including alimentary canal, respiratory tract, excretory system tubes, reproductive system tubes and ducts, body cavity tubes, etc. Local administration or application of the drug(s) affords concentrated delivery of said

2007-01-19 14:24:09

drug(s), achieving tissue levels in target tissues not otherwise obtainable through other administration route.

Means for local drug(s) delivery to hollow tubes can be by physical delivery of the drug(s) either internally or externally to the hollow tube. Local drug(s) delivery includes catheter delivery systems, local injection devices or systems or indwelling devices. Such devices or systems would include, but not be limited to, stents, coated stents, endolumenal sleeves, stent-grafts, liposomes, controlled release matrices, polymeric endolumenal paving, or other endovascular devices, embolic delivery particles, cell targeting such as affinity based delivery, internal patches around the hollow tube, external patches around the hollow tube, hollow tube cuff, external paving, external stent sleeves, and the like. See, Eccleston et al. (1995) Interventional Cardiology Monitor 1:33-40-41 and Slepian, N.J. (1996) Intervente. Cardiol. 1:103-116, or Regar E, Sianos G, Serruys PW. Stent development and local drug delivery. Br Med Bull 2001;59:227-48 which disclosures are herein incorporated by reference.

By "biocompatible" is meant a material which elicits no or minimal negative tissue reaction including e.g. thrombus formation and/or inflammation.

Delivery or application of the drug(s) can occur using stents or sleeves or sheathes. An intraluminal stent composed of or coated with a polymer or other biocompatible materials, e.g. porous ceramic, e.g. nanoporous ceramic, into which the drug(s) has been impregnated or incorporated can be used. Such stents can be biodegradable or can be made of metal or alloy, e.g. Ni and Ti, or another stable substance when intended for permanent use. The drug(s) may also be entrapped into the metal of the stent or graft body which has been modified to contain micropores or channels. Also lumenal and/or ablumenal coating or external sleeve made of polymer or other biocompatible materials, e.g. as disclosed above, that contain the drug(s) can also be used for local delivery.

Stents are commonly used as a tubular structure left inside the lumen of a duct to relieve an obstruction. They may be inserted into the duct lumen in a non-expanded form and are then expanded autonomously (self-expanding stents) or with the aid of a second device in situ, e.g. a catheter-mounted angioplasty balloon which is inflated within the stenosed vessel or body passageway in order to shear and disrupt the obstructions associated with the wall components of the vessel and to obtain an enlarged lumen.

For example, the drug(s) may be incorporated into or affixed to the stent in a number of ways and utilizing any biocompatible materials; it may be incorporated into e.g. a polymer or

2007-10-19 14:24:09

2007-10-19 2:43:09
a polymeric matrix and sprayed onto the outer surface of the stent. A mixture of the drug(s) and the polymeric material may be prepared in a solvent or a mixture of solvents and applied to the surfaces of the stents also by dip-coating, brush coating and/or dip/spin coating, the solvent (s) being allowed to evaporate to leave a film with entrapped drug(s). In the case of stents where the drug(s) is delivered from micropores, struts or channels, a solution of a polymer may additionally be applied as an outlayer to control the drug(s) release; alternatively, the drug may be comprised in the micropores, struts or channels and the adjunct may be incorporated in the outlayer, or vice versa. The drug may also be affixed in an inner layer of the stent and the adjunct in an outer layer, or vice versa. The drug(s) may also be attached by a covalent bond, e.g. esters, amides or anhydrides, to the stent surface, involving chemical derivatization. The drug(s) may also be incorporated into a biocompatible porous ceramic coating, e.g. a nanoporous ceramic coating.

Examples of polymeric materials include biocompatible degradable materials, e.g. lactone-based polyesters or copolyesters, e.g. polylactide; polylactide-glycolide; polycaprolactone-glycolide; polyorthoesters; polyanhydrides; polyaminoacids; polysaccharides; polyphosphazenes; poly(ether-ester) copolymers, e.g. PEO-PLLA, or mixtures thereof; and biocompatible non-degrading materials, e.g. polydimethylsiloxane; poly(ethylene-vinylacetate); acrylate based polymers or copolymers, e.g. polybutylmethacrylate, poly(hydroxyethyl methylmethacrylate); polyvinyl pyrrolidinone; fluorinated polymers such as polytetrafluoroethylene; cellulose esters.

When a polymeric matrix is used, it may comprise 2 layers, e.g. a base layer in which the drug(s) is/are incorporated, e.g. ethylene-co-vinylacetate and polybutylmethacrylate, and a top coat, e.g. polybutylmethacrylate, which is drug(s)-free and acts as a diffusion-control of the drug(s). Alternatively, the drug may be comprised in the base layer and the adjunct may be incorporated in the outlayer, or vice versa. Total thickness of the polymeric matrix may be from about 1 to 20 μ or greater.

According to the method of the invention or in the device or system of the invention, the drug(s) may elute passively, actively or under activation, e.g. light-activation.

The drug(s) elutes from the polymeric material or the stent over time and enters the surrounding tissue, e.g. up to ca. 1 month to 1 year. The local delivery according to the present invention allows for high concentration of the drug(s) at the disease site with low concentration of circulating compound. The amount of drug(s) used for local delivery applications will vary depending on the compounds used, the condition to be treated and the

desired effect. For purposes of the invention, a therapeutically effective amount will be administered. By therapeutically effective amount is intended an amount sufficient to inhibit cellular proliferation and resulting in the prevention and treatment of the disease state. Specifically, for the prevention or treatment of restenosis e.g. after revascularization, or antitumor treatment, local delivery may require less compound than systemic administration.

Preferred combinations are those comprising 40-O-(2-hydroxyethyl)-rapamycin or 32-deoxorapamycin in conjunction or association with a compound having antiproliferative properties, e.g. taxol, paclitaxel, docetaxel, an epothilone, a tyrosine kinase inhibitor, a VEGF receptor tyrosine kinase inhibitor, a VEGF receptor inhibitor, a compound binding to VEGF, a compound having anti-inflammatory properties, e.g. a steroid, a cyclooxygenase inhibitor, ASM981, or with a calcineurin inhibitor, e.g. CysA or FK506. A combination of 40-O-(2-hydroxyethyl)-rapamycin or 32-deoxorapamycin with a compound having anti-inflammatory properties has particularly beneficial effects when used in the treatment or prevention of restenosis in diabetic patients.

Utility of the drug(s) may be demonstrated in animal test methods as well as in clinic, for example in accordance with the methods hereinafter described.

A1. Inhibition of late neointimal lesion formation in the 28 day rat carotid artery balloon injury model

Numerous compounds have been shown to inhibit intimal lesion formation at 2 weeks in the rat ballooned carotid model, while only few compounds prove effective at 4 weeks.

Compounds of formula I are tested in the following rat model.

Rats are dosed orally with placebo or a compound of formula I. Daily dosing starts 3 days prior to surgery and continues for 31 days. Rat carotid arteries are balloon injured using a method described by Clowes *et al.* Lab. Invest. 1983;49;208-215. Following sacrifice at 28 days post-balloon injury, carotid arteries are removed and processed for histologic and morphometric evaluation. In this assay the compounds of formula I, e.g. 40-O-(2-hydroxyethyl)-rapamycin, significantly reduce neointimal lesion formation at 28 days following balloon injury when administered at a dose of from 0.5 to 2.0 mg/kg. For example for 40-O-(2-hydroxyethyl)-rapamycin administered at 0.5, 1.0, and 2.0 mg/kg, the percent inhibition is similar at all three doses: inhibition is 31% at the lowest dose (0.5 mg/kg) and 39% at the highest dose (2.0 mg/kg). Compounds of formula I, e.g. 40-O-(2-hydroxyethyl)-rapamycin, have the beneficial effect to inhibit lesions at 4 weeks post-ballooning

2007-10-14 14:00:02

A.2 Inhibition of restenosis at 28 days in the rabbit iliac stent model

603472154.01.1002

A combined angioplasty and stenting procedure is performed in New Zealand White rabbit iliac arteries. Iliac artery balloon injury is performed by inflating a 3.0 x 9.0 mm angioplasty balloon in the mid-portion of the artery followed by "pull-back" of the catheter for 1 balloon length. Balloon injury is repeated 2 times, and a 3.0 x 12 mm stent is deployed at 6 atm for 30 seconds in the iliac artery. Balloon injury and stent placement is then performed on the contralateral iliac artery in the same manner. A post-stent deployment angiogram is performed. All animals receive oral aspirin 40 mg/day daily as anti-platelet therapy and are fed standard low-cholesterol rabbit chow. Twenty-eight days after stenting, animals are anesthetized and euthanized and the arterial tree is perfused at 100 mmHg with lactated Ringer's for several minutes, then perfused with 10% formalin at 100 mmHg for 15 minutes. The vascular section between the distal aorta and the proximal femoral arteries is excised and cleaned of periadventitial tissue. The stented section of artery is embedded in plastic and sections are taken from the proximal, middle, and distal portions of each stent. All sections are stained with hematoxylin-eosin and Movat pentachrome stains. Computerized planimetry is performed to determine the area of the internal elastic lamina (IEL), external elastic lamina (EEL) and lumen. The neointima and neointimal thickness is measured both at and between the stent struts. The vessel area is measured as the area within the EEL. Data are expressed as mean \pm SEM. Statistical analysis of the histologic data is accomplished using analysis of variance (ANOVA) due to the fact that two stented arteries are measured per animal with a mean generated per animal. A $P < 0.05$ is considered statistically significant.

The compound of formula I, e.g. 40-O-(2-hydroxyethyl)-rapamycin, is administered orally by gavage at a loading dose of 1.5 mg/kg one day prior to stenting, then dosed at 0.75 mg/kg/day from the day of stenting until day 27 post-stenting. In this model, the treatment with the compounds of formula I results in a marked reduction in the extent of restenotic lesion formation: for example, the treatment with 40-O-(2-hydroxyethyl)-rapamycin produces a significant ($P < 0.03$) reduction in neointimal thickness (40% reduction), neointimal area (24% reduction), and percent arterial stenosis (26% reduction) with a significant 32% increase in lumen area. There is extensive neointimal formation in placebo-treated animals at 28 days, with the lesions consisting of abundant smooth muscle cells in proteoglycan/collagen matrix and apparent full endothelial healing. In the majority of arterial segments from the animals treated with 40-O-(2-hydroxyethyl)-rapamycin, the intima is well healed, characterized by a compact neointimal consisting of smooth muscle cells and

endothelium both over stent struts and between struts. Scanning electron microscopic analysis shows that stented arteries from the animals treated with 40-O-(2-hydroxyethyl)-rapamycin (n = 4 arteries) was 84% endothelialized.

A.3 Inhibition of restenosis at 14 days in the rat carotid stent model

Male Sprague Dawley rats weighing 250 to 500 mg are housed individually and allowed to acclimate prior to surgery. All animals receive standard rat chow and water ad libitum. Group size is 12 animals per group.

The drug(s) administration is perivascular. A segment of ballooned carotid is encircled with a 1 cm length of silastic tubing (0.25 inch inside diameter, .047 inch outside diameter) to which is attached a catheter which feeds into an osmotic pump containing either compound or vehicle. This delivery system provides continuous, local delivery to the adventitia of the wrapped portion of vessel. Local drug(s) administration ranges between 5 µg and 10 mg, locally per day, depending on the solubility characteristics of the individual compounds.

The left common carotid arteries are denuded of endothelium using a 2F Fogarty catheter as previously described (Prescott *Am. J. Pathol.* (1991) 139:1291-1296, Clowes et al., (1983) *Lab Invest.* 49:327-333). Briefly, rats are anesthetized with ketamine (50 mg/ml) and rompun (10mg/ml) administered intraperitoneally at a dose of 1.5 ml/kg. A midline incision is made in the neck to expose the left external and common carotid arteries. The balloon is inserted into the common carotid artery via the left external branch, inflated with saline, and pulled back three times through the lumen with a rotating motion to ensure maximal endothelial denudation. The catheter is then removed, the external carotid artery is ligated and the wound is closed. Each animal is given an injection of the antibiotic Bacillin (200,000 units/kg) and the analgesic Buprenorphine (0.06 mg/kg) immediately following surgery.

Animals are killed at 14 days post-balloon injury. One half hour before termination blood is collected, centrifuged, and stored at -20°C for analysis of circulating levels of compound. 5% Evans Blue is then injected intravenously to allow discrimination of re-endothelialized areas at the time of histologic processing. Animals are killed by administration of an overdose of ketamine and rompun, the osmotic pumps are recovered and the volume of remaining content is recorded to ensure that pump failure has not occurred.

Carotid arteries are excised and immersion fixed, then transferred to Ringer's solution. Two samples from control blue region of each left carotid artery are imbedded in paraffin. A minimum of six carotid sections, 20 µM apart are cut per animal and stained with Verhoeff

150347254-011002

Elastic stain to produce a modified Verhoff stain. Intimal and medial area measurements are performed with a computerized imaging system. The intimal lesion area and the medial area are determined by measurement of the internal elastic lamina, the external elastic lamina and the vessel/lumen interface.

In this assay, 40-O-(2-hydroxyethyl)-rapamycin reduces neointimal lesion formation at 14 days post ballooning when administered locally as disclosed above at a dose of 10 to 200 μ g/day. Similar good results are obtained when 40-O-(2-hydroxyethyl)-rapamycin is administered in conjunction with dexamethasone (10-250 μ g/day) or a tyrosine kinase inhibitor.

The following example is illustrative of the invention without limiting it.

A stent is weighed and then mounted for coating. While the stent is rotating, a solution of polylactide glycolide, 0.75 mg/ml of 40-O-(2-hydroxyethyl)-rapamycin, 0.0015 mg/ml 2,6-di-tert.-butyl-4-methylphenol and 1 mg/ml tyrosine kinase inhibitor dissolved in a mixture of methanol and tetrahydrofuran, is sprayed onto it. The coated stent is removed from the spray and allowed to air-dry. After a final weighing the amount of coating on the stent is determined.


2007-07-19 14:26:14

CLAIMS

1. A method for preventing or treating smooth muscle cell proliferation and migration in hollow tubes, or increased cell proliferation or decreased apoptosis or increased matrix deposition in a mammal in need thereof, comprising local administration of a therapeutically effective amount of a rapamycin derivative having mTOR inhibiting properties.
2. A method for the treatment of intimal thickening in vessel walls comprising the controlled delivery from a catheter-based device or an intraluminal medical device of a therapeutically effective amount of a rapamycin derivative having mTOR inhibiting properties.
3. A method according to claim 1 or 2 wherein the rapamycin derivative having mTOR inhibiting properties is administered or delivered in conjunction with one or more other active ingredients selected from a calcineurin inhibitor, an EDG-Receptor agonist, an anti-inflammatory agent, an anti-thrombotic or anti-coagulant agent, an antiproliferative agent, a microtubule stabilizing or destabilizing agent, a tyrosine kinase inhibitor, a compound or antibody which inhibits the PDGF receptor tyrosine kinase or a compound which binds to PDGF or reduces expression of the PDGF receptor, a compound or antibody which inhibits the EGF receptor tyrosine kinase or a compound which binds to EGF or reduces expression of the EGF receptor, a compound or antibody which inhibits the VEGF receptor tyrosine kinase or a VEGF receptor or a compound which binds to VEGF, a statin, a compound, protein, growth factor or compound stimulating growth factor production that will enhance endothelial regrowth of the luminal endothelium, a matrix metalloproteinase inhibitor, a modulator (i.e. antagonists or agonists) of kinases, or a compound stimulating the release of (NO) or a NO donor.

2007-10-19 14:24:09

4. A method according to claim 1 wherein the rapamycin derivative having mTOR inhibiting properties is a compound of formula I



wherein

R₁ is CH₃ or C₃₋₆alkynyl,

R₂ is H, -CH₂-CH₂-OH or 3-hydroxy-2-(hydroxymethyl)-2-methyl-propanoyl, and

X is =O, (H,H) or (H,OH)

provided that R_2 is other than H when X is =O and R_1 is CH_3 .

5. A method according to claim 1 wherein the rapamycin derivative having mTOR inhibiting properties is 40-O-(2-hydroxyethyl)-rapamycin.

6. A drug delivery device or system comprising a) a medical device adapted for local application or administration in hollow tubes, e.g. a catheter-based delivery device or an intraluminal medical device, and b) a therapeutic dosage of a rapamycin derivative having mTOR inhibiting properties being releasably affixed to the medical device.

7. A device according to claim 6 comprising b) a therapeutic dosage of a rapamycin derivative having mTOR inhibiting properties in conjunction with a therapeutic dosage of one or more other active ingredients, each being releasably affixed to the medical device and the other active ingredient being selected from a calcineurin inhibitor, an EDG-Receptor agonist, an anti-inflammatory agent, an anti- thrombotic or anti-coagulant agent, an antiproliferative agent, a microtubule stabilizing or destabilizing agent, a tyrosine kinase inhibitor, a compound or antibody which inhibits the PDGF receptor tyrosine kinase or a compound which binds to PDGF or reduces expression of the PDGF receptor, a compound or antibody which inhibits the EGF receptor tyrosine kinase or a compound which binds to EGF or reduces expression of the EGF receptor, a compound or antibody which inhibits the VEGF receptor tyrosine kinase or a VEGF receptor or a compound which binds to VEGF, a statin, a

8. A device according to claim 6 wherein the rapamycin derivative having mTOR Inhibiting properties is a compound of formula I



R₁ is CH₃ or C₃₋₆alkynyl,

R_2 is H, $-CH_2-CH_2-OH$ or 3-hydroxy-2-(hydroxymethyl)-2-methyl-propanoyl, and

X is =O, (H,H) or (H,OH)

provided that R_2 is other than H when X is =O and R_1 is CH_3 .

9. A device according to claim 6 wherein the rapamycin derivative having mTOR inhibiting properties is 40-O-(2-hydroxyethyl)-rapamycin.

10. A device according to claim 6 comprising b) a therapeutic dosage of rapamycin in conjunction with a therapeutic dosage of one or more other active ingredients, each being releasably affixed to the medical device and the other active ingredient being selected from a calcineurin inhibitor, an EDG-Receptor agonist, an anti-inflammatory agent, a microtubule stabilizing or destabilizing agent, a compound or antibody which inhibits the PDGF receptor tyrosine kinase or a compound which binds to PDGF or reduces expression of the PDGF receptor, a compound or antibody which inhibits the EGF receptor tyrosine kinase or a compound which binds to EGF or reduces expression of the EGF receptor, a compound or antibody which inhibits the VEGF receptor tyrosine kinase or a VEGF receptor or a compound which binds to VEGF, a statin, a compound, protein, growth factor or compound stimulating growth factor production that will enhance endothelial regrowth of the luminal

endothelium, a matrix metalloproteinase inhibitor, an inhibitor of a modulator (i.e. antagonists or agonists) of kinases, and a compound stimulating the release of (NO) or a NO donor.

11. A method according to claim 1 or 2 wherein the administration or delivery is intravascular, intranasal, intrabronchial, interperitoneal or esophageal.

12. A method for preventing or treating smooth muscle cell proliferation and migration in hollow tubes, or increased cell proliferation or decreased apoptosis or increased matrix deposition in a mammal in need thereof, comprising local administration of a therapeutically effective amount of rapamycin in conjunction with a therapeutic dosage of one or more other active ingredients, each being releasably affixed to the medical device and the other active ingredient being selected from a calcineurin inhibitor, an EDG-Receptor agonist, an anti-inflammatory agent, a microtubule stabilizing or destabilizing agent, a compound or antibody which inhibits the PDGF receptor tyrosine kinase or a compound which binds to PDGF or reduces expression of the PDGF receptor, a compound or antibody which inhibits the EGF receptor tyrosine kinase or a compound which binds to EGF or reduces expression of the EGF receptor, a compound or antibody which inhibits the VEGF receptor tyrosine kinase or a VEGF receptor or a compound which binds to VEGF, a statin, a compound, protein, growth factor or compound stimulating growth factor production that will enhance endothelial regrowth of the luminal endothelium, a matrix metalloproteinase inhibitor, an inhibitor of a modulator (i.e. antagonists or agonists) of kinases, and a compound stimulating the release of (NO) or a NO donor.

13. A method for the treatment of intimal thickening in vessel walls comprising the controlled delivery from a catheter-based device or an intraluminal medical device of a therapeutically effective amount of rapamycin in conjunction with a therapeutic dosage of one or more other active ingredients, each being releasably affixed to the medical device and the other active ingredient being selected from a calcineurin inhibitor, an EDG-Receptor agonist, an anti-inflammatory agent, a microtubule stabilizing or destabilizing agent, a compound or antibody which inhibits the PDGF receptor tyrosine kinase or a compound which binds to PDGF or reduces expression of the PDGF receptor, a compound or antibody which inhibits the EGF receptor tyrosine kinase or a compound which binds to EGF or reduces expression of the EGF receptor, a compound or antibody which inhibits the VEGF receptor tyrosine kinase or a VEGF receptor or a compound which binds to VEGF, a statin, a compound, protein, growth factor or compound stimulating growth factor production that will enhance endothelial regrowth of the luminal endothelium, a matrix metalloproteinase

2007-01-01 10:24:54

inhibitor, an inhibitor of a modulator (i.e. antagonists or agonists) of kinases, and a compound stimulating the release of (NO) or a NO donor.

14. A method according to claim 1 or 2 wherein the administration or delivery is made using a catheter delivery system, a local injection device, an indwelling device, a stent, a coated stent, a sleeve, a stent-graft, polymeric endoluminal paving or a controlled release matrix.

15. A method according to claim 1 wherein the rapamycin derivative having mTOR inhibiting properties is administered from a stent or from a coating applied to a stent.

16. A method according to claim 2 wherein the rapamycin derivative having mTOR inhibiting properties is delivered from a stent or from a coating applied to a stent.

17. A method according to claim 1 or 12 for the treatment of stenosis, restenosis or inflammation.

18. A method according to claim 2 or 13 for the treatment of stenosis, restenosis or inflammation.

19. A device according to claim 6 which is a catheter delivery system, a local injection device, an indwelling device, a stent, a stent-graft or a sleeve.

20. A device according to claim 6 which is a coated stent.

21. A device according to claim 7 or 10 which is a catheter delivery system, a local injection device, an indwelling device, a stent, a stent-graft or a sleeve.

22. A device according to claim 7 or 10 which is a coated stent.

2007-09-24

Abstract

The invention relates to the local administration of rapamycin or a derivative thereof, optionally in conjunction with one or more other active ingredients, and a device adapted for such local administration.

50347264:011002